



Redetermination of the crystal structure of dimethylbis[2,4-pentanedionato(1-)-κ²O²,O⁴]-tin(IV)

Hans Reuter* and Martin Reichelt

Institute of Chemistry of New Materials, University of Osnabrück, Barbarastrasse 7, 49069 Osnabrück, Germany.

*Correspondence e-mail: hreuter@uos.de

Received 18 February 2017

Accepted 27 February 2017


Edited by M. Weil, Vienna University of Technology, Austria

Keywords: crystal structure; redetermination; organotin compound; acetylacetonate.**CCDC reference:** 1534819**Supporting information:** this article has supporting information at journals.iucr.org/e

The redetermination of the title compound, [Sn(CH₃)₂(C₅H₇O₂)₂] or SnMe₂(acac)₂, from CCD data recorded at 100 K basically confirms the previous study based on integrated film data recorded at room temperature [Miller & Schlemper (1972). *Inorg. Chem.* **12**, 677–681], but reveals a remarkable shrinkage of the *a* axis [7.12 (1) > 6.7694 (4) Å]. The molecule belongs to point group *C*_i with the Sn^{IV} atom on a centre of inversion. The Sn^{IV} atom shows a slightly distorted octahedral coordination sphere with the methyl groups in *trans* positions and a Sn–C bond length of 2.115 (2) Å which may serve as a standard value for an Sn–CH₃ bond of an octahedrally coordinated Sn^{IV} atom. The Sn–O bonds involving the two carbonyl groups of the acetylacetonate ligand are of equal length [2.180 (1) and 2.183 (1) Å], as are the C=O [1.273 (1) and 1.274 (1) Å] and C–C bond lengths [1.393 (2) and 1.400 (2) Å]. The acetylacetonate ligand deviates considerably from planarity, with a dihedral angle of 5.57 (9)° between the least-squares planes of the two acetone moieties. The four O atoms of the two symmetry-related acetylacetonate ligands are arranged in a nearly quadratic rectangle. Weak C–H···O interactions consolidate the crystal packing.

1. Chemical context


The crystal structure of the title compound, [Sn(CH₃)₂(C₅H₇O₂)₂] or SnMe₂(acac)₂, was determined in the early 1970s at room temperature by visual estimation of film data and refined to a final conventional *R* value of 0.079 (Miller & Schlemper, 1972). All bond lengths and angles of the original study seem chemically reasonable but accuracy suffers from the limited precision of that kind of data collection. As SnMe₂(acac)₂ serves as a reference for all diorganotin(IV) diacetylacetonates and bis-1,3-diketonates in general, we decided to redetermine its structure from CCD data recorded at low temperature. Moreover, the title compound is an excellent candidate for the determination of the Sn–C_{Me} bond length as another reference in case the Sn^{IV} atom is in a well-defined octahedral coordination. The precise measurement of this Sn–C distance therefore should supplement the observations of Britton (2006) who found a significant change in Sn–C bond lengths depending on the organic moiety attached to an Sn^{IV} atom.



2. Structural commentary

The redetermination of the crystal structure of the title compound confirms the former results obtained by Miller & Schlemper (1973) with respect to the chosen space group

($P2_1/n$) and the constitution of the asymmetric unit comprising half a formula unit with the Sn atom at a crystallographic centre of inversion [Wyckhoff symbol: b]. As we performed the X-ray measurement at 100 K, the unit-cell volume is somewhat smaller in comparison with the original room-temperature data which is mainly caused by a considerable change of the a axis from 7.12 (1) to 6.7694 (4) Å while changes of all other lattice parameters [$b_{\text{original}} = 13.87$ (2), $c_{\text{original}} = 7.69$ (1) Å, $\beta_{\text{original}} = 104.7$ (2)°] show a normal temperature-dependent shrinkage.



The Sn atom is octahedrally coordinated with the two methyl groups in a *trans* position (Fig. 1, Table 1). The Sn—C bond shows a length of 2.115 (2) Å and is significantly shorter than the value [2.14 (2) Å] obtained by Miller & Schlemper (1972). Because of the higher coordination number, the value differs to some extent from the value of 2.099 (2) Å for Sn with a coordination number of four as observed in dimethyldithiocyanato tin(IV) (Britton, 2006). The two bonds between the Sn atom and the two different O atoms of the acetylacetonate ligand are of equal length [2.180 (1)/2.183 (1) Å]. In accordance with the almost symmetrical bonding of the acetylacetonate ligand to tin, C—O [1.273 (2)/1.274 (2) Å] and C—C [1.393 (2)/1.400 (2) Å] bonds of the 1,3-diketone skeleton are of equal length. Although these values are typical for a delocalized π system, the atoms in question show a significant deviation from planarity at the central C2 atom, resulting in a dihedral angle of 5.57 (9)° between the least-squares planes defined by O1/C1/C2/C3 [deviations from planarity: 0.003 (1), -0.007 (1), 0.002 (1), 0.002 (1) Å,




Figure 1

The molecular structure of the title compound, showing the atom-labeling scheme of the asymmetric unit. Displacement ellipsoids are drawn at the 50% probability level.

Table 1
Selected geometric parameters (Å, °).

Sn1—C6	2.115 (2)	O1—C1	1.274 (2)
Sn1—C6 ⁱ	2.115 (2)	O2—C3	1.273 (2)
Sn1—O1	2.180 (1)	C1—C2	1.393 (2)
Sn1—O1 ⁱ	2.180 (1)	C2—C3	1.400 (2)
Sn1—O2	2.183 (1)	C3—C5	1.499 (2)
Sn1—O2 ⁱ	2.183 (1)	C1—C4	1.505 (2)
C6 ⁱ —Sn1—C6	180.0	C1—O1—Sn1	125.61 (9)
C6—Sn1—O1	90.20 (5)	O1—C1—C2	126.2 (1)
C6—Sn1—O2	90.41 (5)	C1—C2—C3	128.4 (1)
O1—Sn1—O2	85.99 (4)	C3—O2—Sn1	125.68 (9)

Symmetry code: (i) $-x + 1, -y + 1, -z$.

respectively] and O2/C3/C2/C4 [deviations from planarity: 0.002 (1), -0.004 (1), 0.002 (1), 0.001 (2) Å, respectively] (Fig. 2). Moreover, all bond angles in the six-membered chelate ring are considerably larger [125.61 (9)° at O1, 126.2 (1)° at C1, 128.4 (1)° at C2, 126.07 (13)° at C3, 125.68 (9)° at O2] than expected for sp^2 -hybridized atoms, with exception of the bond angle at Sn1 that amounts to 85.99 (4)°. The four O atoms of the two symmetry-related acetylacetonate ligands around the Sn atom form a planar rectangle with similar edge lengths [O1···O2 = 2.975 (1)/O1···O1 = 3.191 (1) Å], and almost right angles [89.9 (1)° at O1 and 90.1 (1)° at O2]. This plane is nearly perpendicular to the axis through the two methyl groups [deviation: 0.44 (2)°] but constitutes a dihedral angle of 10.2 (1)° with the least-




Figure 2

Twisting of the acetylacetonate ligand at atom C2 with respect to the least-squares planes (green dashed lines) O1/C1/C2/C5 and O2/C3/C2/C4 in a view parallel to these planes. Non-H atoms are shown as displacement ellipsoids at the 50% probability level.




Figure 3

Orientation of the acetylacetonate ligand (least-squares plane through both carbonyl groups) in relation to the plane defined by the four O atoms coordinating to the Sn^{IV} atom. The view is parallel to these planes (green lines). Non-H atoms are shown as displacement ellipsoids at the 50% probability level.




Figure 4
Predominant O...H—C contacts (blue dotted lines) of O atoms with the methyl H atoms of the acetylacetonate groups of neighbouring molecules. The central molecule is drawn in space-filling mode, while neighbouring molecules are drawn in the stick-model mode visualizing the delocalized π system of the acetylacetonate ligands.

squares plane through the two carbonyl groups of the acetylacetonate ligand [deviations from planarity: O1 = 0.006 (1), C1 = -0.007 (1), O2 = -0.006 (1) Å, C3 = 0.007 (1) Å] (Fig. 3).

3. Supramolecular features

In the absence of classical H-donor groups, intermolecular interactions are restricted to van der Waals and weak O...H—C interactions. The most prominent ones are associated with the methyl hydrogen atoms H42 and H51 of the acetylacetonate ligand as they interact simultaneously with both oxygen atoms of neighbouring molecules [C42...O2ⁱ = 2.906 Å, C42...O1ⁱⁱ = 2.852 Å, C51...O2ⁱⁱⁱ = 2.797 Å, H51...O1^{iv} = 2.850 Å; symmetry codes: (i) $x, y, 1 + z$; (ii) $1 - x, 1 - y, 1 - z$; (iii) $\frac{1}{2} + x, \frac{1}{2} - y, \frac{1}{2} + z$; (iv) $\frac{3}{2} - x, -\frac{1}{2} + y, \frac{1}{2} + z$] (Fig. 4). These interactions are completed by a third O...H—C contact of similar length [H2...O2^v = 2.893 Å, H43...O1^v = 2.992 Å, symmetry code: (v) $2 - x, 1 - y, 1 - z$] for each oxygen atom. In summary, the intermolecular contacts result in a columnar arrangement of the molecules parallel to the a axis (Fig. 5).




Figure 5
Columnar arrangement of the molecules along the a axis.

Table 2
Experimental details.

Crystal data	
Chemical formula	[Sn(CH ₃) ₂ (C ₅ H ₇ O ₂) ₂]
M_r	346.97
Crystal system, space group	Monoclinic, $P2_1/n$
Temperature (K)	100
a, b, c (Å)	6.7693 (4), 13.8357 (7), 7.6661 (4)
β (°)	104.709 (2)
V (Å ³)	694.46 (7)
Z	2
Radiation type	Mo $K\alpha$
μ (mm ⁻¹)	1.84
Crystal size (mm)	0.29 × 0.23 × 0.16
Data collection	
Diffractometer	Bruker APEXII CCD
Absorption correction	Multi-scan (SADABS; Bruker, 2009)
T_{\min} , T_{\max}	0.616, 0.760
No. of measured, independent and observed [$I > 2\sigma(I)$] reflections	21644, 1672, 1456
R_{int}	0.024
($\sin \theta/\lambda$) _{max} (Å ⁻¹)	0.660
Refinement	
$R[F^2 > 2\sigma(F^2)]$, $wR(F^2)$, S	0.013, 0.034, 1.11
No. of reflections	1672
No. of parameters	84
$\Delta\rho_{\text{max}}$, $\Delta\rho_{\text{min}}$ (e Å ⁻³)	0.37, -0.27

Computer programs: APEX2 and SAINT (Bruker, 2009), SHELXS97 and SHELXTL (Sheldrick, 2008), SHELXL2014 (Sheldrick, 2015), DIAMOND (Brandenburg, 2006) and Mercury (Macrae *et al.*, 2008).

4. Synthesis and crystallization

The synthesis of the title compound by refluxing a suspension of dimethyltin oxide, Me₂SnO, in acetylacetonate, acacH, for several hours followed the procedure and experimental details described by McGrady & Tobias (1965). Single crystals for X-ray diffraction were grown from toluene solution. A suitable single crystal was selected under a polarization microscope and mounted on a 50 μm MicroMesh MiTeGen Micromount™ using FOMBLIN Y perfluoropolyether (LVAC 16/6, Aldrich). The crystals are stable in air.

Spectroscopic data: ¹H NMR [CDCl₃, TMS, δ (ppm)], nJ [Hz]: $\delta(\text{CH}_3\text{—Sn}) = 0.58$, $^2J(^1\text{H—}^{119/117}\text{Sn}) = 100.6/97.1$; $\delta(\text{CH}_3)_{\text{acac}} = 1.96$; $\delta(\text{CH})_{\text{acac}} = 5.31$ (Lockhart & Manders, 1986; Otera *et al.*, 1981); ¹³C NMR [CDCl₃, TMS, δ (ppm)], nJ [Hz]: $\delta(\text{CH}_3\text{—Sn}) = 7.75$, $^1J(^{13}\text{C—}^{119/117}\text{Sn}) = 973.7/930.4$, $\delta(\text{CH}_3)_{\text{acac}} = 27.94$, $\delta(\text{CH})_{\text{acac}} = 100.09$, $\delta(\text{C=O})_{\text{acac}} = 190.75$ (Lockhart & Manders, 1986; Otera *et al.*, 1981); IR [ATR, ν (cm⁻¹): 3010 w, 2920 w, 1562 s, 1512 s, 1436 m, 1361 s, bd, 1256 m, 1203 m, 1015 m, 925 m, 803 m, 781 m, 655 m, 572 m, 552 m (McGrady & Tobias, 1965); Raman [ν (cm⁻¹): 3092 w, 2999 w, 2920 s, 2708 w, 1574 w, 1427 w, 1366 m, 1263 m, 1206 m, 1194 m, 1021 w, 927 m, 668 m, 567 m, 512 s, 415 m, 220 m, 130 m 94 m, 68 m (McGrady & Tobias, 1965).

5. Refinement details

Crystal data, data collection and structure refinement details are summarized in Table 2. All H atoms were clearly identified

in difference Fourier syntheses but were refined assuming idealized geometries and allowed to ride on the carbon atoms with 0.98 Å (–CH₃), and 0.95 Å (–CH–) and with $U_{\text{iso}}(\text{H}) = 1.2$ and $1.5U_{\text{eq}}(\text{C})$, respectively.

Acknowledgements

We thank the Deutsche Forschungsgemeinschaft and the Government of Lower-Saxony for funding the diffractometer and acknowledge support by Deutsche Forschungsgemeinschaft (DFG) and Open Access Publishing Fund of Osnabrück University.

References

- Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Britton, D. (2006). *Acta Cryst.* **C62**, m93–m94.
- Bruker (2009). *APEX2*, *SADABS* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Lockhart, T. P. & Manders, W. F. (1986). *Inorg. Chem.* **25**, 892–895.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- McGrady, M. M. & Tobias, R. S. (1965). *J. Am. Chem. Soc.* **87**, 1909–1916.
- Miller, G. A. & Schlemper, E. O. (1973). *Inorg. Chem.* **12**, 677–681.
- Otera, J., Hinoishi, T., Kawabe, Y. & Okawara, R. (1981). *Chem. Lett.* **10**, 273–274.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.

supporting information

Acta Cryst. (2017). E73, 472-475 [https://doi.org/10.1107/S2056989017003206]

Redetermination of the crystal structure of dimethylbis[2,4-pentanedionato(1-)- κ^2O^2, O^4]tin(IV)

Hans Reuter and Martin Reichelt

Computing details

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINTE* (Bruker, 2009); data reduction: *SAINTE* (Bruker, 2009); program(s) used to solve structure: *SHELXS97* (Sheldrick 2008); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2015); molecular graphics: *DIAMOND* (Brandenburg, 2006) and *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

Dimethylbis[2,4-pentanedionato(1-)- κ^2O^2, O^4]tin(IV)

Crystal data

[Sn(CH₃)₂(C₅H₇O₂)₂]

$M_r = 346.97$

Monoclinic, $P2_1/n$

$a = 6.7693$ (4) Å

$b = 13.8357$ (7) Å

$c = 7.6661$ (4) Å

$\beta = 104.709$ (2)°

$V = 694.46$ (7) Å³

$Z = 2$

$F(000) = 348$

$D_x = 1.659$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 9878 reflections

$\theta = 2.9$ – 28.5 °

$\mu = 1.84$ mm⁻¹

$T = 100$ K

Prism, colourless

$0.29 \times 0.23 \times 0.16$ mm

Data collection

Bruker APEXII CCD
diffractometer

φ and ω scans

Absorption correction: multi-scan
(*SADABS*; Bruker, 2009)

$T_{\min} = 0.616$, $T_{\max} = 0.760$

21644 measured reflections

1672 independent reflections

1456 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.024$

$\theta_{\max} = 28.0$ °, $\theta_{\min} = 3.0$ °

$h = -8 \rightarrow 8$

$k = -18 \rightarrow 18$

$l = -10 \rightarrow 10$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.013$

$wR(F^2) = 0.034$

$S = 1.11$

1672 reflections

84 parameters

0 restraints

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters not defined?

$w = 1/[\sigma^2(F_o^2) + (0.0135P)^2 + 0.383P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.37$ e Å⁻³

$\Delta\rho_{\min} = -0.27$ e Å⁻³

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Sn1	0.5000	0.5000	0.0000	0.01302 (5)
O1	0.64915 (18)	0.48470 (8)	0.28616 (14)	0.0219 (2)
C1	0.7415 (2)	0.40942 (11)	0.36115 (18)	0.0190 (3)
C2	0.7883 (2)	0.32773 (11)	0.2732 (2)	0.0209 (3)
H2	0.8493	0.2758	0.3489	0.055 (3)*
C3	0.7571 (2)	0.31231 (10)	0.0877 (2)	0.0166 (3)
O2	0.66423 (16)	0.36896 (7)	−0.03729 (13)	0.0183 (2)
C4	0.8070 (3)	0.41338 (14)	0.5638 (2)	0.0314 (4)
H41	0.8498	0.3489	0.6114	0.055 (3)*
H42	0.6924	0.4353	0.6102	0.055 (3)*
H43	0.9214	0.4585	0.6018	0.055 (3)*
C5	0.8410 (3)	0.22215 (11)	0.0252 (2)	0.0272 (3)
H51	0.9046	0.1820	0.1298	0.055 (3)*
H52	0.9435	0.2394	−0.0397	0.055 (3)*
H53	0.7300	0.1860	−0.0553	0.055 (3)*
C6	0.7456 (2)	0.58584 (11)	−0.0348 (2)	0.0236 (3)
H61	0.8232	0.6103	0.0825	0.053 (4)*
H62	0.6921	0.6403	−0.1145	0.053 (4)*
H63	0.8353	0.5466	−0.0887	0.053 (4)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Sn1	0.01680 (7)	0.01050 (7)	0.01066 (7)	0.00215 (5)	0.00147 (5)	−0.00056 (5)
O1	0.0300 (6)	0.0211 (6)	0.0130 (5)	−0.0013 (4)	0.0026 (4)	−0.0031 (4)
C1	0.0147 (6)	0.0287 (8)	0.0123 (6)	−0.0080 (6)	0.0010 (5)	0.0042 (6)
C2	0.0182 (7)	0.0235 (8)	0.0200 (7)	0.0039 (6)	0.0031 (6)	0.0110 (6)
C3	0.0139 (6)	0.0134 (6)	0.0236 (7)	0.0007 (5)	0.0067 (5)	0.0038 (6)
O2	0.0252 (5)	0.0144 (5)	0.0148 (5)	0.0057 (4)	0.0039 (4)	0.0001 (4)
C4	0.0312 (8)	0.0485 (11)	0.0116 (7)	−0.0168 (8)	0.0001 (6)	0.0044 (7)
C5	0.0273 (8)	0.0173 (7)	0.0403 (10)	0.0081 (6)	0.0143 (7)	0.0045 (7)
C6	0.0228 (7)	0.0191 (7)	0.0284 (8)	−0.0012 (6)	0.0058 (6)	0.0009 (6)

Geometric parameters (\AA , $^\circ$)

Sn1—C6	2.115 (2)	C3—C5	1.499 (2)
Sn1—C6 ⁱ	2.115 (2)	C1—C4	1.505 (2)
Sn1—O1	2.180 (1)	C4—H41	0.9800
Sn1—O1 ⁱ	2.180 (1)	C4—H42	0.9800

Sn1—O2	2.183 (1)	C4—H43	0.9800
Sn1—O2 ⁱ	2.183 (1)	C5—H51	0.9800
O1—C1	1.274 (2)	C5—H52	0.9800
O2—C3	1.273 (2)	C5—H53	0.9800
C1—C2	1.393 (2)	C6—H61	0.9800
C2—C3	1.400 (2)	C6—H62	0.9800
C2—H2	0.9500	C6—H63	0.9800
C6 ⁱ —Sn1—C6	180.0	O2—C3—C2	126.07 (13)
C6 ⁱ —Sn1—O1 ⁱ	90.20 (5)	O2—C3—C5	115.27 (13)
C6—Sn1—O1 ⁱ	89.80 (5)	C2—C3—C5	118.66 (13)
C6 ⁱ —Sn1—O1	89.80 (5)	C3—O2—Sn1	125.68 (9)
C6—Sn1—O1	90.20 (5)	C1—C4—H41	109.5
O1 ⁱ —Sn1—O1	180.0	C1—C4—H42	109.5
C6 ⁱ —Sn1—O2	89.59 (5)	H41—C4—H42	109.5
C6—Sn1—O2	90.41 (5)	C1—C4—H43	109.5
O1 ⁱ —Sn1—O2	94.01 (4)	H41—C4—H43	109.5
O1—Sn1—O2	85.99 (4)	H42—C4—H43	109.5
C6 ⁱ —Sn1—O2 ⁱ	90.41 (5)	C3—C5—H51	109.5
C6—Sn1—O2 ⁱ	89.59 (5)	C3—C5—H52	109.5
O1 ⁱ —Sn1—O2 ⁱ	85.99 (4)	H51—C5—H52	109.5
O1—Sn1—O2 ⁱ	94.01 (4)	C3—C5—H53	109.5
O2—Sn1—O2 ⁱ	180.0	H51—C5—H53	109.5
C1—O1—Sn1	125.61 (9)	H52—C5—H53	109.5
O1—C1—C2	126.2 (1)	Sn1—C6—H61	109.5
O1—C1—C4	114.7 (2)	Sn1—C6—H62	109.5
C2—C1—C4	119.1 (1)	H61—C6—H62	109.5
C1—C2—C3	128.4 (1)	Sn1—C6—H63	109.5
C1—C2—H2	115.8	H61—C6—H63	109.5
C3—C2—H2	115.8	H62—C6—H63	109.5

Symmetry code: (i) $-x+1, -y+1, -z$.